

Diff-Stat™

THE MOST POWERFUL FRONT LINE DEFENSE AGAINST
ANTIBIOTIC ASSOCIATED DIARRHEA (AAD) & C-DIFF



Synbiotic Chewable Tablets & Powder Paks

Serving Size: 1 Chewable Tablet or 1 Powder Pak

- *Saccharomyces boulardii* 7.5 billion CFU
- *Lactobacillus Sporogenes* 1 billion CFU
(*Bacillus coagulans*)
- *Fructo-oligosaccharides (FOS)* 500 mg

Indications

Recommended Dose per serving (1 chewable tablet/1 powder pak)

- | | |
|------------------------------------|---------------|
| • AAD & C-diff prophylaxis | 1 time daily |
| • AAD | 2 times daily |
| • C-diff | 2 times daily |
| • Multiple C-diff relapses | 3 times daily |
| • Maintain intestinal tract health | 1 time daily |

**Diff-Stat™ contains 3 clinically proven ingredients
that help combat the effects of antibiotics on
intestinal microbiota.**



MEDICAL NUTRITION USA, Inc.
TRANSLATING SCIENCE INTO SOLUTIONS

Englewood, New Jersey 07631 (800) 221-0308 (201)-569-1188 (in NJ) fax: (201)-569-3224
www.diff-stat.info

Diff-Stat™

POLICY & PROCEDURE

Policy:

The policy of this facility is to assure optimal health and nutrition for all Residents/patients at all times and to help prevent diarrhea, AAD, & C-diff.

Procedure:

Residents/patients who are ordered a broad spectrum antibiotic should take the recommended dose at the start of antibiotic therapy and continue taking at least 2 weeks after completing antibiotics.

Residents/patients who are not currently on antibiotics but are at risk for AAD & C-diff should take the recommended dose for at least 30 days.

Directions for oral consumption:

Diff-Stat Chewable Tablets: Dissolve in mouth or chew.

Diff-Stat Powder Pak (unflavored): Dissolve with 30 ml juice and consume immediately. *

Directions for Diff-Stat Powder Pak through a naso-gastric or stomach tube:

1. Open Diff-Stat Powder Pak away from residents/patients with a central venous catheter (CVC). **
2. Mix Diff-Stat powder with 30 ml of sterile water and administer through naso-gastric or stomach tube immediately. *
3. Flush with 30 ml of sterile water.

Directions for Diff-Stat Powder Pak for residents/patients with a CVC:

1. Open Diff-Stat pak in another room from residents/patients with a central venous catheter (CVC). **
2. Mix Diff-Stat powder with 30 ml of sterile water and place in a syringe. *
3. Change gloves.
4. Administer orally or through naso-gastric/ stomach tube.
5. Flush tube with 30 ml of sterile water.

**Diff-Stat is only viable for 30 minutes in solution.*

*** Saccharomyces boulardii spores could be aerosolized or inadvertently introduced into a CVC line when the pak is open. These spores can float in the air and potentially contaminate the CVC line, therefore it is important to open and mix in another room.*

CAUTION:

Diff-Stat™ should be taken 1 hr before or 2 hrs after antifungal medication.

Do not use if you have a yeast allergy.